

CHEMNITZ

Quantum dynamics for random models: why, what and how GAMM AGUQ Workshop TU Dortmund, 2018

### Quantum dynamics for random models: why, what and how

Peter Stollmann

GAMM AGUQ Workshop TU Dortmund, 2018

13.3.2018



TECHNISCHE UNIVERSITÄT CHEMNITZ

· 13.3.2018 · Peter Stollmann

 < □ > < □ > < □ > < ≥ > < ≥ > < ≥ >
 ≥ < ○ ○</td>

 http://www.tu-chemnitz.de/mathematik/analysis



- 1. The Anderson model ► WHY?
- 2. Quantum dynamics ► WHAT?
- 3. Random models ► WHAT?
- 4. Methods ► HOW?
- 5. References

▲ロト ▲母 ト ▲目 ト ▲目 ト 三目 - のへで





In the celebrated article [4] "Absence of Diffusion in Certain Random Lattices". Phys. Rev. 109 (5): 1492 -1505 from 1958, P.W. Anderson proposed a mathematical model to explain the phase transition from insulator to metal in disordered solids.

Ξ





#### from Anderson's Nobel lecture, [3]: The original Anderson model

· 13.3.2018 · Peter Stollmann

Ξ



Random model for solids: Hopping term plus random onsite potential. In our terminology:

$$H_{\omega}\psi(x) = -\Delta_{disc}\psi(x) + V_{\omega}(x)\psi(x),$$

defines a random operator. The ingredients: x - site on a "lattice"  $\Delta_{disc}$  - Hopping terms  $V_{\omega}$  - random potential

> Need probability distributions not averages: no real atom is an average atom

▲ロト ▲団ト ▲臣ト ▲臣ト 三臣 - のへで

Anderson proved: nontransport theorem, **localization**, and suggested transition, now called Anderson transition.



The Anderson model

Why?

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─ のへで



Quantum dynamics The Schrödinger equation

$$\dot{\psi}(t) = -i(-\Delta + V)\psi(t), \psi(0) = \psi_0,$$



· 13.3.2018 · Peter Stollmann

・ロト・(アト・モト・モト) モークロート
http://www.tu-chemnitz.de/mathematik/analysis



▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

▶ If the spectral measure  $\rho_{\eta_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

▶ If the spectral measure  $\rho_{\eta_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

But what does it look like?

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

▶ If the spectral measure  $\rho_{\eta_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

#### But: what does it look like?

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

▶ If the spectral measure  $\rho_{\eta_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$

· 13.3.2018 · Peter Stollmann



 $\psi(t) = e^{-iHt}\psi_0.$ 

#### But what does it look like? To this end one studies the spectral resolution of H.

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

▶ If the spectral measure  $\rho_{\eta_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$

3



 $\psi(t) = e^{-iHt}\psi_0.$ 

But: what does it look like? To this end one studies the spectral resolution of *H*.

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

One speaks of a **bound state**.

▶ If the spectral measure  $\rho_{\psi_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

But: what does it look like? To this end one studies the spectral resolution of *H*.

• If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

One speaks of a **bound state**.

▶ If the spectral measure  $\rho_{\psi_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

But: what does it look like?

To this end one studies the spectral resolution of H.

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

#### One speaks of a **bound state**.

• If the spectral measure  $\rho_{\psi_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

But: what does it look like?

To this end one studies the spectral resolution of H.

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

One speaks of a **bound state**.

 $\blacktriangleright~$  If the spectral measure  $\rho^H_{\psi_0}$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

But: what does it look like?

To this end one studies the spectral resolution of H.

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

One speaks of a **bound state**.

 $\blacktriangleright~$  If the spectral measure  $\rho^H_{\psi_0}$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

But: what does it look like?

To this end one studies the spectral resolution of H.

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

One speaks of a **bound state**.

 $\blacktriangleright~$  If the spectral measure  $\rho^H_{\psi_0}$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$



 $\psi(t) = e^{-iHt}\psi_0.$ 

But: what does it look like?

To this end one studies the spectral resolution of H.

▶ If  $\psi_0$  is an eigenvector of H with eigenvalue  $E_0 \Rightarrow$ 

$$\psi(t) = e^{-iE_0 t} \psi_0.$$

One speaks of a **bound state**.

▶ If the spectral measure  $\rho_{\psi_0}^H$  of  $\psi_0$  w.r.t H is continuous,

$$\Rightarrow \lim_{T \to \infty} \frac{1}{T} \int_{-T}^{T} \|\chi_B \psi(t)\|^2 dt = 0 \text{ for compact } B.$$

One speaks of a scattering state.

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─ のへで



- Atomic models are described by a negative potential V that decays rapidely enough near infinity: one gets bound states for negative energies and scattering
- ►
- $\triangleright$  P.W. Anderson proposed a new paradigm for disordered solids in dimension  $\geq 3$ :



- Atomic models are described by a negative potential V that decays rapidely enough near infinity:one gets bound states for negative energies and scattering
- ►
- $\triangleright$  P.W. Anderson proposed a new paradigm for disordered solids in dimension  $\geq 3$ :



- Atomic models are described by a negative potential V that decays rapidely enough near infinity: one gets bound states for negative energies and scattering states for positive energies.
- ►.
- $\triangleright$  P.W. Anderson proposed a new paradigm for disordered solids in dimension  $\geq 3$ :

3



- ► Atomic models are described by a negative potential V that decays rapidely enough near infinity:one gets bound states for negative energies and scattering states for positive energies.
- Solid states with perfect order are described by a periodic V ⇒ H = −△ + V has only scattering states.
- **•** P.W. Anderson proposed a new paradigm for disordered solids in dimension  $\geq 3$ :



- ► Atomic models are described by a negative potential V that decays rapidely enough near infinity:one gets bound states for negative energies and scattering states for positive energies.
- Solid states with perfect order are described by a periodic V ⇒ H = −∆ + V has only scattering states.
- **P**.W. Anderson proposed a new paradigm for disordered solids in dimension  $\geq 3$ :



- Atomic models are described by a negative potential V that decays rapidely enough near infinity: one gets bound states for negative energies and scattering states for positive energies.
- Solid states with perfect order are described by a periodic  $V \Rightarrow H = -\Delta + V$ has only scattering states.
- ▶ P.W. Anderson proposed a new paradigm for disordered solids in dimension ≥ 3:

Da C



- ► Atomic models are described by a negative potential V that decays rapidely enough near infinity:one gets bound states for negative energies and scattering states for positive energies.
- ► Solid states with perfect order are described by a periodic  $V \Rightarrow H = -\Delta + V$  has only scattering states.
- P.W. Anderson proposed a new paradigm for disordered solids in dimension  $\geq$  3:

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─ のへで



# Once translated into the language of spectral theory there is a transition from a **localized phase** that exhibits pure point spectrum (= only bound states = no transport)

to a

delocalized phase with absolutely continuous spectrum

(= scattering states = transport)

The dynamical properties of solutions to the Schrödinger equation

 $\dot{\psi}(t) = -i(-\Delta + V)\psi(t), \psi(0) = \psi_0,$ 

depend drastically on the energy of  $\psi_0$ !



#### Once translated into the language of spectral theory there is a transition from a **localized phase** that exhibits pure point spectrum

#### to a

#### delocalized phase with absolutely continuous spectrum

(= scattering states = transport)

$$\dot{\psi}(t) = -i(-\Delta + V)\psi(t), \psi(0) = \psi_0,$$

3



Once translated into the language of spectral theory there is a transition from a **localized phase** that exhibits pure point spectrum

(= only bound states = no transport)

to a

delocalized phase with absolutely continuous spectrum

(= scattering states = transport)

The dynamical properties of solutions to the Schrödinger equation

 $\dot{\psi}(t) = -i(-\Delta + V)\psi(t), \psi(0) = \psi_0,$ 

depend drastically on the energy of  $\psi_0!$ 

▲□▶ ▲圖▶ ▲≣▶ ▲≣▶ = 第二 の�?



#### However, for genuine random models, there is no rigorous proof for the existence of a transition or even of the appearance of spectral components other than pure point, so far.



However, for genuine random models, there is no rigorous proof for the existence of a transition or even of the appearance of spectral components other than pure point, so far.

This is guite a strange situation: the unperturbed problem exhibits extended states and purely a.c. spectrum but for the perturbed one can prove the opposite spectral behavior only.

3



$$H(\omega) = -\Delta + V_{\omega} \text{ in } L^2(\mathbb{R}^d)$$

E.g.,  $\Omega = \mathbb{R}^{\mathbb{Z}^d}$ ,  $\mathbb{P} = \mu^{\mathbb{Z}^d}$  a probability space describing independent, identically distributed coupling constants

$$V_{\omega} = \sum_{k \in \mathbb{Z}^d} \omega_k \cdot v(\cdot - k)$$

sometimes also called alloy type models. Localization has been proven for such models under additional technical hypotheses on  $\mu$  and v.





$$H(\omega) = -\Delta + V_{\omega} \text{ in } L^2(\mathbb{R}^d)$$

with impurities located according to a Poisson process

$$V_{\omega} = \sum_{k \in \mathbb{N}} v(\cdot - X_k(\omega))$$



Localization was shown for such models in [8]

· 13.3.2018 · Peter Stollmann

http://www.tu-chemnitz.de/mathematik/analysis

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─ のへで



Here the operators are given by

$$H(\omega) = -\nabla \mathbf{a}_{\omega} \nabla \operatorname{in} L^2(\mathbb{R}^d)$$

where the leading order coefficient is a randomly chosen matrix, independently in different space regions, more precisely

$$\mathbf{a}_{\omega} = \mathbf{a}_{per} + \sum_{k \in \mathbb{Z}^d} \omega_k \cdot \chi(\cdot - k),$$

and  $\omega = (\omega_k)_{k \in \mathbb{Z}^d} \in S^{\mathbb{Z}^d}$  is picked w.r.t. a product measure, see [6] and [12].

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─ のへで

#### A model studied by Kleespies and P.S.



#### picture taken from [9]

· 13.3.2018 · Peter Stollmann

http://www.tu-chemnitz.de/mathematik/analysis

E

(日) (四) (王) (王) (王)



Much more complicated are the models studied by Borisov and Veselic 2013, [5], where the curvature is random.

One main technical difficulty: missing monotonicity in the dependence on the random variables. In that respect these models share technical features with random displacement models, coming next:



$$H(\omega) = -\Delta + V_{\omega} \text{ in } L^2(\mathbb{R}^d)$$

#### with impurities randomly displaced from their lattice position

$$V_{\omega} = \sum_{k \in \mathbb{Z}^d} v(\cdot - k - \omega_k)$$



FIGURE 1. A typical configuration

#### ... taken from [10].

· 13.3.2018 · Peter Stollmann

17/22

▲ロト ▲母 ト ▲目 ト ▲目 ト 三目 - のへで





Starting from a given graph (e.g.,  $\mathbb{Z}^2$ ) edges are deleted at random and independently; the random operator is the Laplacian on the resulting subgraph, see e.g. [11].

Ξ

- ► New methods are required, traditional perturbation theory fails!
- Statistics of eigenvalues;
- decay properties of eigenfunctions;
- ... for finite volume restrictions of random operators

Two approaches:

- ► Fractional moment method: see Aizenman and Molchanov 1993, [1], for the start and the monograph [2] by Aizenman and Warzel for a recent account.
- Multi-scale analysis: see Fröhlich and Spencer 1983, [7], for the start and the monograph [13] for an extended proof.

▲□▶ ▲□▶ ▲臣▶ ▲臣▶ 三臣 - のへの



Is an induction process proving "good decay properties" of resolvents on boxes  $C_L$  with high probability. With a little bit of oversimplification:

-							
G	G	G	B	G	G	G	G
B	G	G	G	B	G	G	B
G	G	G	G	G	G	G	G
G	G	G	G	G	G	G	G
G	G	G	G	G	G	ß	G
G	G	B	G	G	G	G	G
G	G	G	B	G	G	G	G
G	G	G	G	G	G	B	G

- ► Box of sidelength *L* decomposed into boxes of sidelength  $\ell << L$ .
- Small box is good if resolvent decays exponentially with rate γ<sub>ℓ</sub> > 0, that happens with probability p<sub>ℓ</sub>.
- ► Decay for the large box is controlled by decay for "most" small boxes, giving a decay rate γ<sub>L</sub> < γ<sub>ℓ</sub> > 0 with probability p<sub>L</sub> > p<sub>ℓ</sub>.
- Use independence in the latter step!



Two main ingredients:

• Nonresonance, Wegner estimate: For all  $E \in \mathbb{R}$ ,

 $\mathbb{P}\left(\operatorname{dist}(E, \sigma(H_{\omega} \restriction C_{\ell})) \leq \epsilon\right) \leq C\ell^{d}\epsilon,$ 

where  $H_{\omega} \upharpoonright C_{\ell}$  denotes a selfadjoint restriction of the random operator to the box  $C_{\ell}$ .

► Initial length scale estimates allow to start the induction procedure. For *E* in a certain energy regime, the probability that a box is good (resolvent at energy *E*) is high enough. Often proved in terms of Lifshitz tail estimates.

▲□▶ ▲圖▶ ▲≣▶ ▲≣▶ = 第二 の�?



Methods Multiscale analysis

## Thank you for your attention

· 13.3.2018 · Peter Stollmann

イロト イヨト イヨト イヨト http://www.tu-chemnitz.de/mathematik/analysis

## References CONTRACTOR MARKETAR

- Localization at large disorder and at extreme energies: an elementary derivation. *Comm. Math. Phys.*, 157(2):245–278, 1993.
- Michael Aizenman and Simone Warzel. Random operators, volume 168 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2015. Disorder effects on quantum spectra and dynamics.

#### P. W. Anderson.

Local moments and localized states. *Rev. Mod. Phys.*, 50:191–201, Apr 1978.



#### P.W. Anderson. Absence of diffusion in certain random lattices. *Phys. Reviews*, 109(5):1492–1505, 1958.

Denis Borisov and Ivan Veselić.
 Low lying eigenvalues of randomly curved quantum waveguides.
 J. Funct. Anal., 265(11):2877–2909, 2013.



- Alexander Figotin and Abel Klein. Localization of classical waves. II. Electromagnetic waves. *Comm. Math. Phys.*, 184(2):411–441, 1997.
- Jürg Fröhlich and Thomas Spencer. Absence of diffusion in the Anderson tight binding model for large disorder or low energy. *Comm. Math. Phys.*, 88(2):151–184, 1983.
- François Germinet, Peter D. Hislop, and Abel Klein. Localization for Schrödinger operators with Poisson random potential. J. Eur. Math. Soc. (JEMS), 9(3):577–607, 2007.
- Frank Kleespies and Peter Stollmann. Lifshitz asymptotics and localization for random quantum waveguides. *Rev. Math. Phys.*, 12(10):1345–1365, 2000.
- Frédéric Klopp, Michael Loss, Shu Nakamura, and Günter Stolz. Localization for the random displacement model. Duke Math. J., 161(4):587–621, 2012.

3



- Reza Samavat, Peter Stollmann, and Ivan Veselić. Lifshitz asymptotics for percolation Hamiltonians. Bull. Lond. Math. Soc., 46(6):1113-1125, 2014.

#### Peter Stollmann.

Localization for random perturbations of anisotropic periodic media. Israel J. Math., 107:125-139, 1998.

#### 

Peter Stollmann.

Caught by disorder, volume 20 of Progress in Mathematical Physics. Birkhäuser Boston, Inc., Boston, MA, 2001. Bound states in random media